

Abundant molecular oxygen in the coma of comet 67P/Churyumov–Gerasimenko

A. Bieler^{1,2}, K. Altwegg^{2,3}, H. Balsiger², A. Bar-Nun⁴, J.-J. Berthelier⁵, P. Bochslers², C. Briois⁶, U. Calmonte², M. Combi¹, J. De Keyser⁷, E. F. van Dishoeck⁸, B. Fiethe⁹, S. A. Fuselier¹⁰, S. Gasc², T. I. Gombosi¹, K. C. Hansen¹, M. Hässig^{2,10}, A. Jäckel², E. Kopp², A. Korth¹¹, L. Le Roy³, U. Mall¹¹, R. Maggiori⁷, B. Marty¹², O. Mousis¹³, T. Owen¹⁴, H. Rème^{15,16}, M. Rubin², T. Sémon², C.-Y. Tzou², J. H. Waite¹⁰, C. Walsh⁸ & P. Würz^{2,3}

The composition of the neutral gas comas of most comets is dominated by H₂O, CO and CO₂, typically comprising as much as 95 per cent of the total gas density¹. In addition, cometary comas have been found to contain a rich array of other molecules, including sulfuric compounds and complex hydrocarbons. Molecular oxygen (O₂), however, despite its detection on other icy bodies such as the moons of Jupiter and Saturn^{2,3}, has remained undetected in cometary comas. Here we report *in situ* measurement of O₂ in the coma of comet 67P/Churyumov–Gerasimenko, with local abundances ranging from one per cent to ten per cent relative to H₂O and with a mean value of 3.80 ± 0.85 per cent. Our observations indicate that the O₂/H₂O ratio is isotropic in the coma and does not change systematically with heliocentric distance. This suggests that primordial O₂ was incorporated into the nucleus during the comet's formation, which is unexpected given the low upper limits from remote sensing observations⁴. Current Solar System formation models do not predict conditions that would allow this to occur.

Measurements of the coma of 67P/Churyumov–Gerasimenko (hereafter 67P) were made between September 2014 and March 2015 with the ROSINA-DFMS mass spectrometer⁵ on board the Rosetta spacecraft. For the present study, we analysed 3,193 mass spectra taken in this time period. Because of the high mass resolving power and sensitivity of ROSINA-DFMS, it was possible to differentiate unambiguously between the three main species present in the narrow mass range centred at mass-to-charge (m/z) ratio 32 Da/e, namely, molecular oxygen (O₂), atomic sulfur (S) and methanol (CH₃OH); such differentiation has not been achieved by previous *in situ* or remote sensing measurements at comets. Figure 1 shows several DFMS measurements centred at the O₂ peak. The green and orange lines show data taken before the close encounter with 67P. Only minor signatures from the tenuous neutral gas atmosphere of the Rosetta spacecraft can be identified, and even after long thruster firing manoeuvres, which use N₂O₄ as an oxidizer, the contamination of the O₂ signal remains small (see the green line in Fig. 1). Measurements taken while orbiting 67P, shown as the light blue, dark blue and purple lines in Fig. 1, show a clear increase of the O₂ signal, indicative of cometary O₂. These three measurements were taken at decreasing distances (r) from the comet nucleus, and follow the predicted $1/r^2$ signal dependence that is expected for conserved cometary species, giving further confidence in our detection.

As previously reported, the local number densities in the coma vary spatially and temporally^{6,7} for different compounds. Figure 2 shows

correlation plots of H₂O with O₂, CO and N₂. Red symbols represent data from 17 to 23 October 2014, for which previously published data on N₂ are available. During this time the spacecraft was in closed orbits at 10 km from the nucleus centre. Black symbols are data from 1 September 2014 to 31 March 2015. For both periods, O₂ clearly shows the strongest correlation with H₂O. While CO shows a high correlation with H₂O from 17 to 23 October 2014, the correlation for the whole data set is fairly low. N₂ shows the weakest correlation with H₂O of all three species. The strong correlation between H₂O and O₂, with a Pearson correlation coefficient of 0.88 (and even 0.97 for the October data), indicates that they are of similar origin in the nucleus and that their release mechanisms are linked, in contrast to CO and




Figure 1 | DFMS mass spectra around 32 Da/e normalized to the spectrum with the largest signal. The black labels indicate the three major species found in the coma of 67P at 32 Da/e. The green labels and green line identify contamination peaks from thruster firings, showing that their contributions to the O₂ peak are very low. The light blue, dark blue and purple lines represent measurements taken at different distances from the comet nucleus.

¹Department of Climate and Space Science and Engineering, University of Michigan, 2455 Hayward Street, Ann Arbor, Michigan 48109, USA. ²Physikalisches Institut, University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland. ³Center for Space and Habitability, University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland. ⁴Department of Geosciences, Tel-Aviv University, Ramat-Aviv, 6997801 Tel-Aviv, Israel. ⁵LATMOS/IPSL-CNRS-UPMC-UVSQ, 4 Avenue de Neptune, F-94100 Saint-Maur, France. ⁶Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), UMR 6115 CNRS – Université d'Orléans, 45071 Orléans, France. ⁷Belgian Institute for Space Aeronomy, BIRA-IASB, Ringlaan 3, B-1180 Brussels, Belgium. ⁸Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands. ⁹Institute of Computer and Network Engineering (IDA), TU Braunschweig, Hans-Sommer-Straße 66, D-38106 Braunschweig, Germany. ¹⁰Space Science and Engineering Division, Southwest Research Institute, 6220 Culebra Road, San Antonio, Texas 78228, USA. ¹¹Max-Planck-Institut für Sonnensystemforschung, Justus-von-Liebig-Weg 3, 37077 Göttingen, Germany. ¹²Centre de Recherches Pérogaphiques et Géochimiques, CRPG-CNRS, Université de Lorraine, 15 rue Notre Dame des Pauvres, BP 20, 54501 Vandoeuvre lès Nancy, France. ¹³Aix Marseille Université, CNRS, LAM (Laboratoire d'Astrophysique de Marseille) UMR 7326, 13388 Marseille, France. ¹⁴Institute for Astronomy, University of Hawaii, Honolulu, Hawaii 96822, USA. ¹⁵Université de Toulouse–UPS–OMP–IRAP, 31400 Toulouse, France. ¹⁶CNRS–IRAP, 9 avenue du Colonel Roche, BP 44346, F-31028 Toulouse Cedex 4, France.




Figure 2 | Correlation between H₂O and O₂, CO and N₂. **a**, H₂O and O₂; **b**, H₂O and CO; **c**, H₂O and N₂. All three panels share a common y axis. Numbers on x and y axes are proportional to number density but in arbitrary units. Red crosses mark a subset of data for which N₂ data are also available. Panel **a** shows the strong correlation between H₂O and O₂, which is

observed for all data. In contrast, the correlation of CO with H₂O (**b**) varies over time, which leads to a low overall correlation between those two species. N₂ has the lowest correlation with H₂O of the compared species for the October data (**c**).

N₂ which have a similar volatility but do not show a strong correlation with H₂O (see Fig. 2 for correlation coefficient values). The O₂/H₂O ratio decreases for high H₂O abundances, which might be caused by surface water ice produced by a cyclic sublimation–condensation process⁸, although the total amount of surface ice is limited⁹.

A plausible mechanism for the strong O₂/H₂O correlation would be the production of O₂ by radiolysis or photolysis of water ice. Here we follow the convention that photolysis refers to ultraviolet photons breaking bonds, whereas radiolysis refers to more energetic photons or fast electrons and ions depositing energy into the ice and ionizing molecules. Creation of sputtered O₂ by radiolysis has been demonstrated in laboratory experiments¹⁰ and is observed for the icy moons of Jupiter—Europa, Ganymede and Callisto^{11–13}—as well as for the rings of Saturn³. Comets are subject to radiolysis over various timescales: (a) over billions of years, while they reside in the Kuiper belt; (b) over the period of a few years once they enter the inner Solar System; and (c) on very short timescales, as for the present radiolysis. In the Kuiper belt, the skin depth for producing O₂ is in the range of metres, although the produced O₂ may diffuse deeper into the porous nucleus. Once a comet begins its residence in the inner Solar System, it loses its surface material to a depth of several metres during each orbit around the Sun, therefore we can safely assume that no O₂ from radiolysis in the Kuiper belt phase remains in 67P at the percentage level. Radiolysis and photolysis by solar wind and ultraviolet radiation in the inner Solar System only affect the top few micrometres of the cometary

surface. Taking account of 67P's continuous mass loss through outgassing, we estimate the actively outgassing surface areas to be lost to a depth of several centimetres over the time from August 2014 to March 2015. If recent production by radiolysis or photolysis (only affecting the top few micrometres) were the source of the measured O₂, our data would show a continuous decrease of the O₂/H₂O ratio over the examined time period as the active surface continues to be shed over that time. Apart from the variations related to H₂O abundance, Fig. 3 shows that we do not observe a systematic change in the O₂/H₂O ratio over several months. Instantaneous creation of the measured O₂ by radiolysis or photolysis seems, overall, unlikely, and would lead to variable O₂ ratios due to different illumination conditions. Given that radiolysis and photolysis, on any of the discussed timescales, do not seem to be plausible production mechanisms, the preferred explanation of our observations is the incorporation of primordial O₂ into the cometary nucleus.

Despite great efforts by remote sensing campaigns, information on primordial O₂ is still limited. Solid O₂ has not yet been detected in interstellar ices, and upper limits for the O₂/H₂O ice ratios of <0.5 and for O₂/CO ratios of <1 are in agreement with our findings, but such high upper limits do not provide useful constraints^{14,15}. Gaseous O₂ has only been detected in two interstellar clouds so far^{4,16,17}, and is generally known to have surprisingly low abundances⁴. Reports of very low upper limits for O₂ in a protostellar envelope suggest the material infalling to the accretion disk is very poor in molecular oxygen¹⁸. This has been ascribed




Figure 3 | O₂/H₂O ratio over several months. There seems to be no systematic increase or decrease of the O₂/H₂O ratio. The variances happen on very short timescales and can be explained by the decrease of the O₂ ratio for high H₂O abundances. It is not fully understood if the higher variability

of the O₂ ratio from October to the end of December 2014 can be attributed to orbital changes of the spacecraft or to physical changes of the cometary nucleus.




Figure 4 | DFMS spectra for some of the common products of radiolysis of water ice. **a–c**, Products are O_2H (seen in **a**), $^{16}\text{O}^{18}\text{O}$ and O_2H_2 (seen in **b**) and O_3 (not seen in **c**). These data were recorded on 20 October 2014 at around 01:00 UTC. With the exception of O_3 (**c**), all the previously mentioned

species are measured and can clearly be identified in the mass spectra of DFMS. The grey area consists of the statistical error N/\sqrt{N} and a 10% uncertainty from the individual pixel gains on the detector (here N is the number of molecules on the detector).

to the high volatility and reactivity of O_2 , in particular the rapid transformation of O and O_2 to H_2O ice on cold grains¹⁹.

However, in one of the two interstellar clouds where O_2 has been detected (the ρ Ophiuchi dense core), the chemically related species HO_2 and H_2O_2 have been measured^{20,21} with gaseous abundance ratios of $\text{HO}_2/\text{O}_2 \approx \text{H}_2\text{O}_2/\text{O}_2 \approx 0.6 \times 10^{-3}$. Interestingly, the abundance ratios determined by DFMS for the coma of 67P are very close to these interstellar values: $\text{HO}_2/\text{O}_2 = (1.9 \pm 0.3) \times 10^{-3}$ and $\text{H}_2\text{O}_2/\text{O}_2 = (0.6 \pm 0.07) \times 10^{-3}$ (see Fig. 4). If these gas-phase abundance ratios reflect those in the cometary ice, this would support the existence of primordial O_2 . The ρ Ophiuchi A core has been suggested to have experienced slightly higher temperatures of around 20–30 K over its lifetime (which is also typical of estimates for the comet-forming conditions in the outer early solar nebula), compared to ~ 10 K for most other dense interstellar clouds^{21,22}. If higher temperatures are indeed needed to produce significant amounts of O_2 , this would indicate that our Solar System was formed from an unusually warm molecular cloud, consistent with the low abundance of N_2 in 67P (ref. 23).

One aspect of the present results that remains unexplained is the high value (a few per cent) of the $\text{O}_2/\text{H}_2\text{O}$ ratio in 67P. Models of gas-grain chemistry in molecular clouds predict $\text{O}_2/\text{H}_2\text{O}$ ratios at least an order of magnitude lower¹⁹. They also over-predict ozone (O_3): we found no evidence for the presence of ozone (see Fig. 4), with an upper limit of 1×10^{-6} relative to water.

An alternative explanation for the presence of O_2 is the incorporation of gaseous O_2 into water ice in the protosolar nebula, and chemical models of protoplanetary disks do in fact show high abundances of gaseous O_2 in the comet forming zone²⁴. Rapid cooling from >100 K to less than 30 K would then be needed to form amorphous water ice with trapped O_2 on dust grains. This could happen when young disks experience increased heating due to accretion bursts onto the star, followed by a rapid drop in temperature as soon as the burst is over. These O_2 rich grains then need to be accreted into larger bodies before further chemical modification occurs.

Finally, we discuss radiolysis of icy grains before accretion. When produced by radiolysis in water ice, O_2 can remain trapped in voids, while hydrogen can diffuse out¹⁰. This prevents the hydrogenation of O_2 , which is otherwise a dominant reaction for the destruction of molecular oxygen, and could lead to increased and stable levels of O_2 in the solid ice²⁵. Incorporation of such icy grains into the comet nucleus would explain the observed strong correlation with H_2O , in contrast to N_2 which is trapped from the gas phase and shows a lower correlation with water. However, O_3 (resulting from O_2 radiolysis) has been

reported to be trapped in Ganymede's surface²⁵, and at such a concentration would just be detectable by DFMS for O_2 levels at a few per cent of H_2O , but no O_3 could be detected. A further consequence would be that these icy grains have been incorporated into the comet mostly unaltered, a process that is much debated, but which has recently been proposed again²⁶ and that would also be in accordance with the measured high D/H ratio in 67P (ref. 27).

We note that our findings do not significantly affect our understanding of the global distribution of elemental O in the interstellar medium, as O_2 in ice with an abundance of a few per cent relative to H_2O accounts only for a small fraction of the total oxygen inventory.

Online Content Methods, along with any additional Extended Data display items and Source Data, are available in the online version of the paper; references unique to these sections appear only in the online paper.

Received 29 June; accepted 18 August 2015.

- Bockelée-Morvan, D., Mumma, M. J. & Weaver, H. A. in *Comets II* (eds Festou, M., Keller, U. H. & Weaver, H. A.) 391–423 (Univ. Arizona Press, 2004).
- Hall, D. T., Strobel, D. F., Feldman, P. D., McGarth, M. A. & Weaver, H. A. Detection of an oxygen atmosphere on Jupiter's moon Europa. *Nature* **373**, 677–679 (1995).
- Johnson, R. E. *et al.* Production, ionization and redistribution of O_2 in Saturn's ring atmosphere. *Icarus* **180**, 393–402 (2006).
- Goldsmith, P. F. *et al.* Herschel measurements of molecular oxygen in Orion. *Astrophys. J.* **737**, 96 (2011).
- Balsiger, H. *et al.* ROSINA — ROSETTA orbiter spectrometer for ion and neutral analysis. *Space Sci. Rev.* **128**, 745–801 (2007).
- Hässig, M. *et al.* Time variability and heterogeneity in the coma of 67P/Churyumov-Gerasimenko. *Science* **347**, aaa0276 (2015).
- Luspai-Kuti, A. *et al.* Composition-dependent outgassing of comet 67P/Churyumov-Gerasimenko from ROSINA/DFMS — implications for nucleus heterogeneity? *Astron. Astrophys.* <http://dx.doi.org/10.1051/0004-6361/201526205> (2015).
- De Sanctis, M. C. *et al.* The diurnal cycle of water ice on comet 67P/Churyumov-Gerasimenko. *Nature* **525**, 500–503 (2015).
- Capaccioni, F. *et al.* The organic-rich surface of comet 67P/Churyumov-Gerasimenko as seen by VIRTIS/Rosetta. *Science* **347**, aaa0628 (2015).
- Brown, W. L. *et al.* Erosion and molecular formation in condensed gas films by electronic energy loss of fast ions. *Nucl. Instrum. Methods* **198**, 1–8 (1982).
- Carlson, R. W. *et al.* Hydrogen peroxide on the surface of Europa. *Science* **283**, 2062–2064 (1999).
- Spencer, J. R., Calvin, W. M. & Person, M. J. Charge-coupled device spectra of the Galilean satellites: molecular oxygen on Ganymede. *J. Geophys. Res.* **100**, 19049–19056 (1995).
- Spencer, J. R. & Calvin, W. M. Condensed O_2 on Europa and Callisto. *Astron. J.* **124**, 3400–3403 (2002).
- Vandenbussche, B. *et al.* Constraints on the abundance of solid O_2 in dense clouds from ISO-SWS and ground-based observations. *Astron. Astrophys.* **346**, L57–L60 (1999).
- Pontoppidan, K. *et al.* A 3–5 μm VLT spectroscopic survey of embedded young low mass stars I. Structure of the CO ice. *Astron. Astrophys.* **408**, 981–1007 (2003).

16. Liseau, R. *et al.* Multi-line detection of O₂ toward ρ Ophiuchi A. *Astron. Astrophys.* **541**, A73 (2012).
17. Larsson, B. *et al.* Molecular oxygen in the Ophiuchi cloud. *Astron. Astrophys.* **466**, 999–1003 (2007).
18. Yildiz, U. A. *et al.* Deep observations of O₂ toward a low-mass protostar with Herschel-HIFI. *Astron. Astrophys.* **558**, A58 (2013).
19. Taquet, V., Ceccarelli, C. & Kahane, C. Multilayer modeling of porous grain surface chemistry. I. The GRAINOBLE model. *Astron. Astrophys.* **538**, A42 (2012).
20. Bergman, P. *et al.* Detection of interstellar hydrogen peroxide. *Astron. Astrophys.* **531**, L8 (2011).
21. Parise, B., Bergman, P. & Du, F. Detection of the hydroperoxyl radical HO₂ toward ρ Ophiuchi A. Additional constraints on the water chemical network. *Astron. Astrophys.* **541**, L11 (2012).
22. Du, F., Parise, B. & Bergman, P. Production of interstellar hydrogen peroxide (H₂O₂) on the surface of dust grains. *Astron. Astrophys.* **538**, A91 (2012).
23. Rubin, M. *et al.* Molecular nitrogen in comet 67P/Churyumov-Gerasimenko indicates a low formation temperature. *Science* **348**, 232–235 (2015).
24. Walsh, C., Nomura, H. & van Dishoeck, E. The molecular composition of the planet-forming regions of protoplanetary disks across the luminosity range. *Astron. Astrophys.* **582**, A88 (2015).
25. Johnson, R. E. & Jesser, W. A. O₂/O₃ microatmospheres in the surface of Ganymede. *Astrophys. J.* **480**, L79–L82 (1997).
26. Cleaves, L. I. *et al.* The ancient heritage of water ice in the solar system. *Science* **345**, 1590–1593 (2014).
27. Altwegg, K. *et al.* 67P/Churyumov-Gerasimenko, a Jupiter family comet with a high D/H ratio. *Science* **347**, <http://dx.doi.org/10.1126/science.1261952> (2015).

Acknowledgements Work at the University of Michigan was funded by NASA contract JPL-1266313. Work at the University of Bern was funded by the State of Bern, the Swiss

National Science Foundation and the European Space Agency PRODEX Program. Work at Max-Planck-Institut für Sonnensystemforschung was funded by the Max-Planck Society and BMWI contract 50QP1302. Work at Southwest Research Institute was supported by subcontract 1496541 from the Jet Propulsion Laboratory. Work at BIRA-IASB was supported by the Belgian Science Policy Office via PRODEX/ROSINA PEA 90020. This work was carried out thanks to the support of the A*MIDEX project (no. ANR-11-IDEX-0001-02) funded by the 'Investissements d'Avenir' French Government programme, managed by the French National Research Agency (ANR). This work was supported by CNES grants at IRAP, LATMOS, LPC2E, UTINAM, CRPG, and by the European Research Council (grant no. 267255 to B.M.). A.B.-N. thanks the Ministry of Science and the Israel Space agency. Work by J.H.W. at Southwest Research Institute was funded by NASA JPL subcontract NAS703001TONM0710889. E.F.v.D. and C.W. are supported by A-ERC grant 291141 CHEMPLAN and an NWO Veni award. We acknowledge here the work of the whole ESA Rosetta team.

Author Contributions A.B. performed data reduction, analysis and wrote the paper; K.A. initialized and edited the paper and contributed to data interpretation; C.B., U.C., M.C., T.J.G., K.C.H., S.G., M.H., A.J., R.M., L.L.R., M.R., C.-Y.T. and T.S. contributed to data analysis and interpretation. A.B.-N. and O.M. contributed to data interpretation relevant to processes in ices. E.F.v.D. and C.W. contributed to data interpretation and writing of sections concerning interstellar oxygen. H.B., J.-J.B., P.B., J.D.K., B.F., S.A.F., A.K., U.M., B.M., T.O., H.R., J.H.W. and P.W. contributed to experiment design, calibration and data interpretation. All authors discussed the results, and commented on and revised the manuscript.

Author Information All ROSINA-DFMS data will be released to the PSA archive of ESA and to the PDS archive of NASA. Reprints and permissions information is available at www.nature.com/reprints. The authors declare no competing financial interests. Readers are welcome to comment on the online version of the paper. Correspondence and requests for materials should be addressed to A.B. (abieler@umich.edu).

METHODS

Data reduction. The integration time for all evaluated spectra is 20 s. A single mass spectrum consists of the signals of 512 individual pixels, which are arranged along the dispersive axis of the mass spectrometer. We take the gain degradation of each individual pixel over time into account by determining its gain with a calibration sequence dedicated to this task every few weeks.


For every recorded mass spectrum, a third-degree polynomial is fitted to the baseline of the mass spectrum and subsequently subtracted from the signal. To avoid contamination in the baseline fit from the measured peaks, the centre of the mass spectrum (with the peaks) is ignored for those fitting purposes. Peaks are fitted with a Gaussian-shape curve. This introduces a systematic underestimation of the absolute magnitude of the signal, as at the 1% level and below the peaks are broader than Gaussian, but cancels out for the reported ratios of O₂/H₂O as both species are underestimated similarly. In a post-processing procedure, outliers are removed. An outlier occurs if one of the following criteria is met: (a) the signal amplitude changes by more than two orders of magnitudes between two measurements; (b) the centre of the fitted peak is not within a given window around the predicted position; (c) the width of the fitted peak curve is wider than a set limit.

For the CO measurements, the fraction contributed to the signal by CO fragments from CO₂ is subtracted. As DFMS measures one mass line at a time in high resolution mode, there are no simultaneous measurements of two species that are separated by more than 1 Da/e. To calculate ratios, the two measurements have to occur within 20 min; if no pair of measurements can be found within 20 min, the data point is ignored. In our error analysis we include uncertainties for the individual gain levels of the detector, the species branching ratios and their relative sensitivities as systematic errors. Individual pixel gains are treated as statistical error, as the peak position on the detector is not constant. A careful analysis confirmed that the O₂/H₂O ratio is independent of the peak centre positions.

Sample size. No statistical methods were used to predetermine sample size.

Spacecraft outgassing background. To clearly identify the measured O₂ as cometary in origin, all non-cometary sources of O₂ must be considered and excluded. The Rosetta spacecraft produces a neutral gas cloud of its own, mainly due to diffusion of volatiles out of spacecraft material and desorption of re-deposited volatiles from the spacecraft. For example, by changing the spacecraft attitude, different spacecraft elements are illuminated by the Sun, which then warm up and release condensed gas. The orange line in Fig. 1 shows the low-level signals from this spacecraft contamination for O₂, S and CH₃OH, referred to as “background”. Those measurements were taken several days before the encounter with 67P. It is not possible to distinguish this background signal from any potential cometary signature with DFMS, but it has been well characterized before the arrival at 67P and is usually orders of magnitude lower than the measured O₂ signals²⁸. To keep the background influence as low as possible, we only considered mass spectra where both the O₂ and H₂O abundances are at least 5 times larger than the corresponding spacecraft contamination. Another potential source of O₂ is the oxidizer, N₂O₄, used by the Rosetta spacecraft during thruster firings. Measurements taken shortly after a large thruster firing manoeuvre from June 2014 (still before arrival at the comet) show minor contaminations around 32 Da/e, but not directly affecting the O₂ peak (see green curve in Fig. 1). Although contamination from thruster firings is small, DFMS measurements are usually performed hours after thruster firings, in order to minimize influence thereof. Finally we can exclude the production of O₂ inside the instrument through a careful review of all oxygen-bearing molecules up to 150 Da/e, which could potentially fragment into O₂ in the DFMS electron impact ion source. Many minor species contain O₂ but these are too low in abundance to account for the large amount of O₂ detected. The remaining possibility is CO₂, which is very abundant in the coma of 67P (ref. 6). However, owing to its molecular structure it only fragments into CO and O, not into O₂ (ref. 29). Finally, we exclude the production of O₂ from H₂O in the instrument. For 81 mass spectra taken from May to the end of June 2014 (before the encounter with 67P) we determine an O₂ abundance of (0.18 ± 0.07)% relative to H₂O, which is a factor of 20 lower than the cometary values.

Coma chemistry. The production of O₂ from atomic oxygen in the neutral gas coma of 67P is possible through the reaction



Owing to the tenuous nature of the coma, only very few collisions are expected within the first 100 km above the nucleus surface³⁰. With increasing activity and therefore denser coma one would expect a higher O₂/H₂O ratio as both reactants (atomic oxygen and OH) are related to H₂O as well as the collision frequency. This is clearly not observed. Production of a significant amount of O₂ from coma chemistry hence seems unfeasible.

Correlation with H₂O. The measured O₂ signal shows a very strong dependence on radial distance (*r*) from the comet. It increases by roughly one order of magnitude when the radial distance from the comet decreases from 30 km to 10 km. This is in agreement with a predicted 1/*r*² dependence of the number density profile of a non-reactive species. Examining the data further, we observe a strong correlation between H₂O and O₂ (see Fig. 2, Pearson correlation coefficient *R* = 0.88) for data from September 2014 to March 2015. This correlation indicates that O₂ and H₂O are both of a similar cometary origin. In contrast, there is no correlation between O₂ and H₂O for measurements taken before the arrival at the comet (*R* = −0.01). The observed temporal variations in the O₂/H₂O ratio are largely due to a nonlinear correlation between H₂O and O₂ for high water densities, where the O₂ ratio drops with increasing H₂O abundance. The correlation similarly supports the ruling out of CO₂ as a source of the O₂, as previous studies^{6,23} and Fig. 2 show a lack of correlation between the abundance of H₂O and other species like CO, CO₂ and N₂.

Radiolysis. The production of O₂ from water ice by radiolysis is the result of several reactions, where initially H, O and OH are produced, followed by subsequent rearrangement to form H₂, HO₂ and H₂O₂ and ultimately O₂ (ref. 2). In a systematic study of the irradiation of pure cubic crystalline water ice at 10 K, it was estimated that comets in the Oort cloud would reach an equilibrium concentration of about 0.6% of H₂O₂ after ~10⁵ years, and that the concentration of O₂ would be significantly lower³¹. As stated by the authors, the relative proportion of O₂ and H₂O₂ in actual astrophysical ices can be different as the presence of H₂O₂ ices may enhance the production of O₂ and vice versa.

Cosmic rays consist of energetic particles (mostly H⁺) that can penetrate inside the cometary nucleus. Their penetration depth depends on their energy. The more energetic a cosmic ray, the deeper it deposits its energy into the cometary nucleus. Only the most energetic components, the Galactic cosmic rays (GCRs), can penetrate deeply into the cometary body. Today, the GCR flux peaks at energies around a few hundreds of MeV. At energies higher than ~1 GeV, the GCR flux drops exponentially, the flux at 10 GeV being approximately 100 times lower than the flux at 1 GeV. GCRs with energies near the GCR maximum of flux typically deposit their energy in the nucleus at depths of the order of metres to tens of metres³². There is some uncertainty on the evolution of the cosmic ray flux over the history of the Solar System related to the occurrence of supernovae which can drastically increase the cosmic ray flux at high energies (up to ~PeV) during relatively short time periods (of the order of 10⁵ years) or to the orbital motion of the Sun and of the Galaxy. There are also uncertainties on the cross-sections, the relative proportion of high-Z cosmic rays, and the role of porosity and defects in the cometary ice³³.

During 67P's stay in the Kuiper belt (~4.5 billion years) GCRs may have produced a significant amount of O₂. This O₂ is produced where the GCR energy is deposited, that is, in the first tens of metres below the comet surface. Once cometary objects enter the inner Solar System they lose (owing to their activity and depending on their perihelion distance) material from the surface in the range of several metres per orbit around the Sun. Therefore, as most products built up during the stay in the Kuiper belt reside in the outer few metres, O₂ produced by radiolysis should be released quickly on the first solar passages. 67P's perihelion distance has been within the orbit of Jupiter for the past 250 years, possibly even for more than 5,000 years, and since a close encounter with Jupiter in 1959, the perihelion distance of 67P has been about 1.3 AU with an orbital period of 6.4 years (ref. 34). Accumulated over the last perihelion passages we can assume that 67P has lost hundreds of metres from its surface.

However, O₂ molecules may have diffused into the comet and penetrated into deeper regions of the cometary nucleus. According to the estimated erosion of the comet, this requires significant diffusion of the O₂ molecules. Such efficient diffusion would dilute the produced O₂, resulting in a lower O₂/H₂O ratio hardly compatible with the elevated amount of O₂ observed by DFMS.

Recent radiolysis. The flux of the low energy component such as solar energetic particles (SEP) is several orders of magnitude higher than the flux of GCRs. The SEP flux peaks at energies around a few tens of MeV and only penetrates the first centimetres of the comet³². Considering the comet erosion rate, the production of O₂ by low energy cosmic rays would have to be very fast to account for the observed amount of O₂, requiring unrealistically high O₂ production rates. Consequently, it appears unlikely that most of the O₂ in 67P has been built up over the past few years.

O₂ production through surface sputtering. It has been shown that sputtering of refractory materials from the cometary surface due to the solar wind is occurring at 67P (ref. 35) and a clear difference between the southern and northern latitudes in the measured abundances of the sputtered species was demonstrated. This apparent spatial difference is explained via the asymmetry of the neutral coma, with higher number densities in the northern hemisphere that is preferentially exposed to the Sun for the time span under study. The solar wind, which is responsible for

the sputtering, is therefore attenuated more efficiently by these denser parts of the coma and thus has limited or no access to the surface. We find that the O_2/H_2O ratio is independent of latitude and relatively constant over a period of 7 months. Furthermore, the major part of the top surface of 67P accessible by the solar wind does not contain any water ice⁹. This suggests that sputtering cannot be the main source of the detected molecular oxygen. Moreover, the sputter yields by solar wind ions are orders of magnitude too low to explain the observed amount of O_2 .

Code availability. The data reduction software was written in the Julia and Python programming languages and is available upon request from A.B.

28. Schläppi, B. *et al.* Influence of spacecraft outgassing on the exploration of tenuous atmospheres with in situ mass spectrometry. *J. Geophys. Res.* **115**, A12313 (2010).
29. Tian, C. & Vidal, C. R. Electron impact dissociative ionization of CO_2 : measurements with a focusing time-of-flight mass spectrometer. *J. Chem. Phys.* **108**, 927 (1998).
30. Fuselier, S. A. *et al.* ROSINA/DFMS and IES observations of 67P: ion-neutral chemistry in the coma of a weakly outgassing comet. *Astron. Astrophys.* <http://dx.doi.org/10.1051/0004-6361/201526210> (2015).
31. Zheng, W., Jewitt, D. & Kaiser, R. I. Formation of hydrogen, oxygen and hydrogen peroxide in electron-irradiated crystalline water ice. *Astrophys. J.* **639**, 534–548 (2006).
32. Cooper, J. F., Christian, E. R. & Johnson, R. E. Heliospheric cosmic ray irradiation of Kuiper belt comets. *Adv. Space Res.* **21**, 1611–1614 (1998).
33. Grieves, G. A. & Orlando, T. M. The importance of pores in the electron stimulated production of D_2 and O_2 in low temperature ice. *Surf. Sci.* **593**, 180–186 (2005).
34. Maquet, L. The recent dynamical history of comet 67P/Churyumov-Gerasimenko. *Astron. Astrophys.* **579**, A78 (2015).
35. Wurz, P. *et al.* Solar wind sputtering of dust on the surface of 67P/Churyumov-Gerasimenko. *Astron. Astrophys.* <http://dx.doi.org/10.1051/0004-6361/201525980> (2015).